

This report provides an update of the African swine fever (ASF) situation, according to the information submitted through the World Animal Health Information System of the World Organisation for Animal Health (OIE-WAHIS) between 20 January and 3 February 2022.

The information included in this report was reported by countries through Immediate notifications (IN), follow-up reports (FUR) and six-monthly reports (SMR). More details on OIE data collection for OIE-listed diseases is available on the OIE website¹.

This report will cover: (1) ASF distribution and the situation in 2020-2022 (based on INs, FURs and SMRs) and (2) the recent updates that occurred during the 2-week period of 20 January – 3 February 2022 (based on INs and FURs).

ASF distribution and the situation in 2020 - 2022 (based on INs, FURs and SMRs)

ASF has traditionally been present in the African continent, where since 2005 the disease has been reported in 32 countries. In 1978, the disease was introduced to the Italian island of Sardinia and has since become endemic. In 2007, the disease was confirmed in the Caucasus region of Georgia. From there, the ASF virus gradually spread to neighboring countries (i.e., Armenia, Azerbaijan, Russia and Belarus) affecting domestic pigs and wild boar. The first occurrence of ASF was reported in the European Union (EU) in 2014 and since then, numerous EU countries have been affected by this devastating pig disease that continues to be reported in 16 countries (during 2020 / 2022). Two European countries have managed to eradicate the disease: Belgium (event resolved in March 2020) and Czech Republic (event resolved in April 2018).

In August 2018, the virus leapt to China (People's Rep. of), which represented the first occurrence of ASF in Asia. Since then, the disease continued to spread in the Region, affecting 16 countries as of 2021.

In September 2019, the first occurrence of ASF in Oceania was reported by Timor-Leste, followed by Papua New Guinea (March 2020). And in July 2021 the disease reappeared in the Americas after an absence of almost 40 years, having been introduced in Dominican Republic and later in Haiti. In January 2022, ASF genotype II was notified on the Italian mainland after around 40 years of absence. Two new countries reported the first occurrence of the disease in the country in January as well: North Macedonia in Europe and Thailand in Asia.

¹ <https://www.oie.int/en/what-we-do/animal-health-and-welfare/disease-data-collection/>

ASF distribution in 2020-2022 (as of 05 February 2022) is shown in Figure 1.




Figure 1. Map of the world displaying the presence of ASF by Administrative divisions (2020 – 05/02/2022)

Summary of the ASF situation by world region (2020-2022)

In total, since January 2020 ASF has been reported as present in five different world regions in 35 countries, affecting more than 1,000,000 pigs and more than 30,000 wild boars (data reported through INs and FURs), with more than 1,700,000 animal losses. Further details, split by world region are included in Table 1.

Table 1: Summary of the number of outbreaks, cases and animal losses caused by ASF in the different world regions since January 2020 (data reported through INs and FURs – these figures cover only epizootic situation while additional information reported through SMR for enzootic situation are not included here because of submission delays).

	Outbreaks		Cases		Losses*
	Domestic pigs	Wild boar	Domestic pigs	Wild boar	Domestic pigs
Africa	154		12,991		20,338
Americas	225		8,642		15,130
Asia	1058	1,519	89,389	1,631	398,862
Europe	3,392	16,913	940,204	28,747	1,326,423
Oceania	4		500		397
Total	4,833	18,432	1,051,726	30,378	1,761,150

*Losses (deaths + animals killed and disposed of): this figure refers to losses in the establishments affected by the outbreaks and it does not include the animals culled in areas around the outbreak for controlling the disease.

Summary of the global situation and recommendations

Since January 2020, 8 countries have reported ASF as a first occurrence in the country, while 12 countries reported its spread to new zones. This highlights a continuous spread of the disease into new countries, and new zones in countries already affected.

As observed in Europe and in some regions of Asia, the transmission of ASF seems to depend largely on the wild boar population density and their interaction with low-biosecurity pig production systems. The good knowledge and management of the wild boar population and a good coordination among the Veterinary Services, wildlife and forestry authorities are required to successfully prevent and control ASF.

OIE Members are recommended to implement strict [biosecurity measures](#) and to strengthen their early disease detection system, in particular where there is evidence of circulation of low virulent strains of African swine fever virus, or transmission in wild suids, and to promptly notify any cases of ASF to the OIE.

Recent updates (20/01/2022 – 03/02/2022)

To describe the current disease situation of ASF, this section covers: (a) a list of new events which started during the 2-week period (reported through INs); (b) information on events that started before the 2-week period but were still ongoing during the period (reported through FURs); (c) new events which started before the 2-week period but were reported through INs during the 2-week period and (d) the geographic distribution of new outbreaks that started during the 2-week period. This information is based on INs and FURs received by the OIE.

New events by world region (reported through INs)

Africa, Asia, Americas, Oceania

No new events reported

Europe

Recurrence in Bulgaria started on 24 January

Recurrence in Russia started on 20 January

Recurrence in Ukraine started on 1 February

First occurrence in a zone in Russia (Bashkortostan administrative division) started on 25 January

On-going events for which there were new outbreaks, by world region (reported through FURs):

Americas, Oceania

No ongoing events updated

Africa

One country updated its ongoing events: South Africa

Asia

One country updated its ongoing events: Thailand

Europe

Five countries updated their ongoing events: Hungary, Italy, Latvia, Romania, Russia




Figure 2: Map of ASF outbreaks which started during 20 January – 3 February 2022 in domestic animals and wildlife. Zoomed views of areas where updates occurred in the last period are provided as well.

New events by world region (reported through INs) which started before the reporting periodAsia, Africa, Americas, Oceania

No new events reported

Europe

Recurrence in Bulgaria started on 12 January

Recurrence in Moldova started on 17 January

Recurrence in Ukraine started on 17 January

Discussion

The events observed in the last 6 months confirm the global threat of ASF, which continues to spread in several regions, affecting territories which have previously not reported the disease, with serious impacts on pig production systems, animal health and welfare, as well as on livelihoods, national food security and international trade.

The detection of ASF in these new territories could pose a risk of spread to neighbouring territories, and the OIE encourages Veterinary Services to maintain vigilance, and implement science-based international standards and guidelines in their national prevention and control programmes.

In particular, an early detection system for ASF could facilitate early reporting and response, limiting the spread of disease. ASF surveillance needs to be adapted to the local epidemiological context, taking into account the presence of low virulent strains that could preclude clinical surveillance. Surveillance programmes should also cover wild and feral suid populations where they are involved in the disease epidemiology. OIE Members should also ensure access to quality laboratory diagnosis for ASF, capable of identifying ASF virus in accordance with the standards in the OIE Terrestrial Manual.

Biosecurity is still the most important and most effective measure available to prevent and control ASF. Rigorous and continuous implementation of biosecurity and maintaining a high level of awareness of ASF among all those in the value chain, as well as maintaining vigilance at borders to prevent the illegal movement of ASF-infected commodities, can prevent the virus from entering pig herds.

The control of ASF requires sustained commitment and resources, and the involvement of all relevant stakeholders. Public-private partnerships are in this regard instrumental in leveraging the respective strengths, knowledge, expertise, and resources of both public and private sector partners to allow ASF control to be achieved more rapidly and efficiently.

OIE Members are also reminded that there is no authorised ASF vaccine with proven effectiveness and safety available in the world. Therefore, any type of ASF vaccine sold on the market is either a fake vaccine or contains poorly attenuated strains of ASFV, which poses serious safety risks and has the potential to spread between pigs, causing chronic disease.

The OIE urges its Members to continue to promptly notify the occurrence of ASF and to share the relevant epidemiological information that can facilitate transparency and assist the global control of ASF.