

INFECTION WITH SARS-COV-2 IN ANIMALS

Last updated in February 2022

Severe Acute Respiratory Syndrome-Coronavirus-2 (SARS-CoV-2) is the pathogenic agent that causes the disease COVID-19. Although genetically closely related viruses have been isolated from *Rhinolophus* bats, the exact source of SARS-CoV-2 and route of introduction into the human population has not been established.

The current COVID-19 pandemic is being sustained through human-to-human transmission. Animal infections with SARS-CoV-2 have been [reported](#) by several countries. A range of mammalian species have proven to be susceptible to infection with SARS-CoV-2, with varying clinical manifestations, either naturally and/or by experimental infection. The major livestock species, such as cattle, small ruminants, poultry and swine, do not seem to be susceptible to natural infection. Further studies are needed to understand if and how different animals could be affected by SARS-CoV-2.

It is important to monitor infections in animals to better understand their epidemiological significance for animal health, biodiversity, and human health. Evidence from risk assessments, epidemiological investigations, and experimental studies indicate that animals do not play a significant role in the spread of SARS-CoV-2.

Infection with SARS-CoV-2 in animals is not included in the [OIE List of Diseases](#). However, consistent with the reporting obligations of Members outlined in Article 1.1.4. of the OIE *Terrestrial Animal Health Code* relating to emerging diseases, the disease should be notified to the OIE through the [OIE's World Animal Health Information System](#) or via [email](#).

The information presented in this technical factsheet reflects the current epidemiological evidence and research done to date and will be updated when additional information is available.

AETIOLOGY

Classification of the causative agent

Coronaviruses (CoVs) are enveloped, positive-sense, single-stranded RNA viruses. SARS-CoV-2 is a *betacoronavirus*, a genus that includes several coronaviruses (SARS-CoV, MERS-CoV, bat SARS-like CoV, and others) isolated from humans, bats, camels, civets, and other animals.

Susceptibility to physical and chemical action

SARS-CoV-2 is inactivated by

- 62–71% ethanol, 0.5% hydrogen peroxide or 0.1% sodium hypochlorite, within 1 minute, or
- 0.05–0.2% benzalkonium chloride or 0.02% chlorhexidine digluconate with less effectivity.

Survival:

Under experimental conditions, SARS-CoV-2 remained viable in the environment after aerosolization for at least 180 minutes. Also, under experimental conditions, it was shown that SARS-CoV-2:

- can persist on surfaces such as plastic, stainless steel or glass for 3-7 days, depending on the initial viral load and environmental conditions, but can be efficiently inactivated by surface disinfection procedures, as listed above
- can persist in cloth, paper, wood, and bank notes for 1-2 days
- was found to remain infectious for 14 days at 4°C, but for only 2 days at 20°C in sewage water.

Epidemiology

Hosts

Although current evidence suggests that SARS-CoV-2 emerged from an animal source, likely a bat, that source has yet to be identified. The pandemic is being driven by human-to-human transmission, which happens through respiratory droplets from coughing, sneezing, and talking, that can remain in the air for some time as aerosols. Genetic sequence data reveal that SARS-CoV-2 is genetically closely related to other coronaviruses circulating in *Rhinolophus* bat (horseshoe bat) populations. To date, there is not enough scientific evidence to identify the source of SARS-CoV-2 or to explain the original route of transmission to humans (which may involve an intermediate host).

Several mammalian species have tested positive for SARS-CoV-2, with infection being introduced to a population as a result of close contact with humans or animals infected with SARS-CoV-2 or in experimental infection studies performed in laboratory settings. More information about susceptible species, transmissibility and clinical signs can be found [here](#). The OIE publishes a monthly situation update containing all events on infection of animals with SARS-CoV-2 reported by its Members through OIE-WAHIS. The latest update can be found [here](#).

Transmission

Information on the routes of transmission of SARS-CoV-2 is growing due to field findings and numerous experimental infection studies. As with other respiratory viruses, SARS-CoV-2 is transmitted between humans and animals and between animals by direct contact (e.g., droplets) and through aerosols that can persist in closed environments for some time. SARS-CoV-2 has been found in secretions from the respiratory tract and in faeces.

Viraemia, incubation and infectious period

In laboratory settings, the incubation period in animals appears to be similar to that seen in humans (i.e., between 2 and 14 days). However, more studies are required to better estimate the mean duration of incubation and the infectious periods for the different susceptible animal species.

Sources of virus

The main source of the virus is respiratory droplets, aerosols, and respiratory secretions, although it is possible to isolate SARS-CoV-2 from faeces of some infected animal species.

Pathogenesis

In laboratory settings, infected animals showed presence of the virus in the respiratory tract and, in some cases, lesions in the trachea and lungs associated with dyspnoea and cough.

Occurrence and impact

There have been reports of companion animals and captive wild animals infected with SARS-CoV-2. With respect to production animals, to date, SARS-CoV-2 is known to have affected mink farms in multiple countries, with varying degrees of morbidity and mortality.

Diagnosis

Knowledge about the susceptibility of different animal species to SARS-CoV-2 infection and clinical signs is, to date, limited.

Clinical diagnosis

Knowledge about clinical disease manifestations in animals is limited and there are a range of clinical manifestations. Current evidence suggests clinical signs may include, but are not limited to, coughing, sneezing, respiratory distress, nasal discharge, ocular discharge, vomiting or diarrhoea, fever, inappetence, and lethargy. As in humans, asymptomatic infections can occur in animals.

Lesions

More studies are needed to systematically categorise the lesions resulting from infection with SARS-CoV-2 in animals. The current knowledge is summarised in the following publication from Meekins et al. (2021) <https://doi.org/10.3390/v13101993>

Differential diagnosis

All other causes for respiratory or digestive illness should be excluded before a tentative diagnosis for infection with SARS-CoV-2 is made. Existence of an epidemiological link with a confirmed infection, in humans or other animals, should be considered when narrowing down the list of differential diagnoses.

Laboratory confirmatory tests are necessary for a final diagnosis.

Laboratory diagnosis

Samples

Depending on the type of test, samples may include single or combinations of nasal, oropharyngeal, and rectal swabs, and blood. Faecal samples may be used in situations where direct sampling is not possible due to risks to the animal or testing staff. Tests should be validated for the purpose, species and matrix to be analysed.

Procedures

Agent identification

- Reverse-transcription polymerase chain reaction (RT-PCR)
- Reverse transcription loop-mediated isothermal amplification (RT-LAMP)
- Virus isolation
- Virus genome sequencing
- Other molecular tests developed for use in humans.

Detection of immune response

- ELISA antibody test
- Virus neutralisation test (VNT)
- Several other tests for antibody detection.

Prevention and control

Biosecurity and hygiene measures are key to preventing transmission of SARS-CoV-2.

People who are suspected or confirmed to be infected with SARS-CoV-2 should restrict contact with mammalian animals, including pets, just like they would with people during their illness.

Animals suspected or confirmed to be infected with SARS-CoV-2 should remain separated from other animals and humans while infected.

Due to their susceptibility, some animal species are being used as models to test vaccines for use in humans.

SARS-CoV-2 vaccines for use in mink and other animal species susceptible to infection with SARS-CoV-2 have been used by several countries in farms and zoos.

References

1. World Health Organization, (consulted on 11/05/2020) <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
2. M. Denis, V. Vanderweerd, R. Verbeeke, A. Laudisoit, L. Wynants, D. Van Der Vliet (2020). COVIPENDIUM: information available to support the development of medical countermeasures and interventions against COVID-19 (Version 2020-05-05). Transdisciplinary Insights. <http://doi.org/10.5281/zenodo.3782325>

3. Questions and Answers on COVID-19, (consulted on 11/05/2020), <https://www.oie.int/en/scientific-expertise/specific-information-and-recommendations/questions-and-answers-on-2019novel-coronavirus/>
4. Considerations for sampling, testing, and reporting of SARS-CoV-2 in animals, (consulted on 11/05/2020), https://www.oie.int/fileadmin/Home/eng/Our_scientific_expertise/docs/pdf/COV-19/Sampling_Testing_and_Reporting_of_SARS-CoV-2_in_animals_final_7May_2020.pdf
5. Cohen J. (2020). From mice to monkeys, animals studied for coronavirus answers. *Science*, Vol. 368, Issue 6488 pp. 221-222 <https://science.sciencemag.org/content/368/6488/221>
6. CDC, Coronavirus Disease 2019 (COVID-19) – pets & other animals (consulted on 29/05/2020) <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/positive-pet.html>
7. Schlottau K., Rissmann M., Graaf A., Schön J., Sehl J., Wylezich C., Höper D., Mettenleiter T.C., Balkema-Buschmann A., Harder T., Grund C., Hoffmann D., Breithaupt A., & Beer M. (2020). SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *The Lancet. Microbe*, 1(5), e218–e225. [https://doi.org/10.1016/S2666-5247\(20\)30089-6](https://doi.org/10.1016/S2666-5247(20)30089-6)
8. Sit T., Brackman C.J., Ip S.M., Tam K., Law P., To E., Yu V., Sims L.D., Tsang D., Chu D., Perera R., Poon L., & Peiris M. (2020). Infection of dogs with SARS-CoV-2. *Nature*, 10.1038/s41586-020-2334-5. Advance online publication. <https://doi.org/10.1038/s41586-020-2334-5>
9. Shi J., Wen, Z., Zhong, G., Yang, H., Wang, C., Huang, B., Liu, R., He, X., Shuai, L., Sun, Z., Zhao, Y., Liu, P., Liang, L., Cui, P., Wang, J., Zhang, X., Guan, Y., Tan, W., Wu, G., Chen, H., Bu, Z. (2020). Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science (New York, N.Y.)*, 368(6494), 1016–1020. <https://doi.org/10.1126/science.abb7015>
10. Muñoz-Fontela C., Dowling W.E., Funnell S.G.P. *et al.* Animal models for COVID-19. *Nature* (2020). <https://doi.org/10.1038/s41586-020-2787-6>
11. Oude Munnink, B. B., Sikkema, R. S., Nieuwenhuijse, D. F., Molenaar, R. J., Munger, E., Molenkamp, R., van der Spek, A., Tolsma, P., Rietveld, A., Brouwer, M., Bouwmeester-Vincken, N., Harders, F., Hakze-van der Honing, R., Wegdam-Blans, M., Bouwstra, R. J., GeurtsvanKessel, C., van der Eijk, A. A., Velkers, F. C., Smit, L., Stegeman, A., ... Koopmans, M. (2021). Transmission of SARS-CoV-2 on mink farms between humans and mink and back to humans. *Science (New York, N.Y.)*, 371(6525), 172–177. <https://doi.org/10.1126/science.abe5901>
12. Aboubakr, H. A., Sharafeldin, T. A., & Goyal, S. M. (2020). Stability of SARS-CoV-2 and other coronaviruses in the environment and on common touch surfaces and the influence of climatic conditions: A review. *Transboundary and emerging diseases*, 10.1111/tbed.13707. Advance online publication. <https://doi.org/10.1111/tbed.13707>
13. Pickering, B. S., Smith, G., Pinette, M. M., Embury-Hyatt, C., Moffat, E., Marszal, P...Lewis, C. E. (2021). Susceptibility of Domestic Swine to Experimental Infection with Severe Acute Respiratory Syndrome Coronavirus 2. *Emerging Infectious Diseases*, 27(1), 104-112. <https://dx.doi.org/10.3201/eid2701.203399>.
14. Anna Z. Mykytyn, Mart M. Lamers, Nisreen M. A. Okba, Tim I. Breugem, Debby Schipper, Petra B. van den Doel, Peter van Run, Geert van Amerongen, Leon de Waal, Marion P. G. Koopmans, Koert J. Stittelaar, Judith M. A. van den Brand & Bart L. Haagmans (2021) Susceptibility of rabbits to SARS-CoV-2, *Emerging Microbes & Infections*, 10:1, 1-7, DOI: 10.1080/22221751.2020.1868951
15. Ulrich, L., Wernike, K., Hoffmann, D., Mettenleiter, T. C., & Beer, M. (2020). Experimental Infection of Cattle with SARS-CoV-2. *Emerging Infectious Diseases*, 26(12), 2979-2981. <https://dx.doi.org/10.3201/eid2612.203799>.
16. Freuling, C. M., Breithaupt, A., Müller, T., Sehl, J., Balkema-Buschmann, A., Rissmann, Mettenleiter, T. C. (2020). Susceptibility of Raccoon Dogs for Experimental SARS-CoV-2 Infection. *Emerging Infectious Diseases*, 26(12), 2982-2985. <https://dx.doi.org/10.3201/eid2612.203733>
17. Cool, K., Gaudreault, N. N., Morozov, I., Trujillo, J. D., Meekins, D. A., McDowell, C., Carossino, M., Bold, D., Kwon, T., Balaraman, V., Madden, D. W., Artiaga, B. L., Pogranichniy, R. M., Sosa, G. R., Henningson, J., Wilson, W. C., Balasuriya, U., García-Sastre, A., & Richt, J. A. (2021). Infection and transmission of SARS-CoV-2 and its alpha variant in pregnant white-tailed deer. *bioRxiv : the preprint server for biology*, 2021.08.15.456341. <https://doi.org/10.1101/2021.08.15.456341>
18. Michelitsch, A., Wernike, K., Ulrich, L., Mettenleiter, T. C., & Beer, M. (2021). SARS-CoV-2 in animals: From potential hosts to animal models. *Advances in Virus Research*, 110, 59–102. <https://doi.org/10.1016/BS.AIVIR.2021.03.004>
19. Singh, D. K., Singh, B., Ganatra, S. R., Gazi, M., Cole, J., Thippeshappa, R., Alfson, K. J., Clemmons, E., Gonzalez, O., Escobedo, R., Lee, T. H., Chatterjee, A., Goetz-Gazi, Y., Sharan, R., Gough, M., Alvarez, C., Blakley, A., Ferdin, J., Bartley, C., Staples, H., ... Kaushal, D. (2021). Responses to acute infection with SARS-

- CoV-2 in the lungs of rhesus macaques, baboons and marmosets. *Nature microbiology*, 6(1), 73–86. <https://doi.org/10.1038/s41564-020-00841-4>
20. Meekins, D. A., Gaudreault, N. N., & Richt, J. A. (2021). Natural and Experimental SARS-CoV-2 Infection in Domestic and Wild Animals. *Viruses*, 13(10), 1993. doi:10.3390/v13101993
 21. Chandler, J. C., Bevins, S. N., Ellis, J. W., Linder, T. J., Tell, R. M., Jenkins-Moore, M., Root, J. J., Lenocho, J. B., Robbe-Austerman, S., DeLiberto, T. J., Gidlewski, T., Kim Torchetti, M., & Shriner, S. A. (2021). SARS-CoV-2 exposure in wild white-tailed deer (*Odocoileus virginianus*). *Proceedings of the National Academy of Sciences of the United States of America*, 118(47), e2114828118. <https://doi.org/10.1073/pnas.2114828118>
 22. Wernike, K. *et al.* (2022) 'Serological screening suggests single SARS-CoV-2 spillover events to cattle', *bioRxiv*, p. 2022.01.17.476608. doi: 10.1101/2022.01.17.476608.
 23. Kuchipudi, S. V. *et al.* (2022) 'Multiple spillovers from humans and onward transmission of SARS-CoV-2 in white-tailed deer', *Proceedings of the National Academy of Sciences*, 119(6), p. e2121644119. doi: 10.1073/PNAS.2121644119.