

## CHAPTER 3.1.5.

# CRIMEAN–CONGO HAEMORRHAGIC FEVER

---

### SUMMARY

Crimean–Congo haemorrhagic fever virus (CCHFV) of the genus *Orthonairovirus* of the family *Nairoviridae* causes a zoonotic disease in many countries of Asia, Africa, the Middle East and south-eastern Europe. As the distribution of CCHFV coincides with the distribution of its main vector, ticks of the genus *Hyalomma*, the spread of infected ticks into new, unaffected areas facilitates the spread of the virus. The virus circulates in a tick–vertebrate–tick cycle, but can also be transmitted horizontally and vertically within the tick population. *Hyalomma* ticks infest a wide spectrum of different wildlife species, e.g. deer and hares, and free-ranging livestock animals, e.g. goat, cattle, and sheep. Many birds are resistant to infection, but ostriches appear to be more susceptible. Viraemia in livestock is short-lived, and of low intensity. These animals play a crucial role in the life cycle of ticks, and in the transmission and amplification of the virus and are, therefore, in the focus of veterinary public health. As animals do not develop clinical signs, CCHFV infections have no effect on the economic burden regarding livestock animal production. In contrast to animals, infections of humans can result in the development of a severe disease, Crimean–Congo haemorrhagic fever (CCHF).

Every year, more than 1000 human CCHF cases are reported with case fatality rates of 5–80% depending on the virus strain and other local factors. The pathogenesis of the disease in humans is not well understood. Most people become infected by tick bites and by crushing infected ticks, but infection is also possible through contact with blood and other body fluids of viraemic animals, for example in slaughterhouses. As CCHFV also has the potential to be transmitted directly from human-to-human, nosocomial outbreaks have been reported.

There is no approved CCHF vaccine available and therapy is restricted to treatment of the symptoms. Health education and information on prevention and behavioural measures are most important in order to enhance public risk perception and, therefore, decrease the probability of infections. Thus the identification of endemic areas is crucial for focused and targeted implementation of public health measures. Serological screening of ruminants allows CCHFV-affected areas to be identified, as antibody prevalence in animals is a good indicator of local virus circulation. Treatment with tick repellents can be quite effective in reducing the tick infestation of animals. To protect laboratory staff, handling of CCHFV infectious materials should only be carried out at an appropriate biocontainment level.

**Detection and identification of agent:** Only a single virus serotype is known to date although sequencing analysis indicates considerable genetic diversity. CCHFV has morphological and physiochemical properties typical of the family *Nairoviridae*. The virus has a single-stranded, negative-sense RNA genome consisting of three segments: L (large), M (medium) and S (small), each of which is contained in a separate nucleocapsid within the virion. The virus can be isolated from serum or plasma samples collected during the febrile or viraemic stage of infection, or from liver of infected animals. Primary isolations are made by inoculation of several tissue cultures, commonly African green monkey kidney (Vero) cells. For identification and characterisation of the virus, conventional and real-time reverse transcription polymerase chain reaction (PCR) can be used. As infections of animals remain clinically unapparent, the likelihood of isolating virus from a viraemic animal is very low.

**Serological tests:** Type-specific antibodies are demonstrable by indirect immunofluorescence test or by IgG-sandwich and IgM-capture enzyme-linked immunosorbent assay. Commercial test systems are available for animal health; in addition a few in-house systems have been published or kits are used replacing the conjugate provided in kit with one that is suitable for the animal species to be screened for CCHFV-specific antibodies.

**Requirements for vaccines:** There is no vaccine available for animals.

## A. INTRODUCTION

Crimean-Congo haemorrhagic fever (CCHF) is a zoonotic disease caused by a primarily tick-borne CCHF virus (CCHFV) of the genus *Orthonairovirus* of the family *Nairoviridae*, order *Bunyavirales*. CCHFV possesses a negative-sense RNA genome consisting of three segments, L (large), M (medium) and S (small) each contained in a separate nucleocapsid within the virion. All orthonairoviruses are believed to be transmitted by either ixodid or argasid ticks, and several are known to be pathogenic to humans, namely CCHF, Dugbe and Nairobi sheep disease viruses (Swanepoel & Burt, 2004; Swanepoel & Paweska, 2011; Whitehouse, 2004). CCHFV can be grown in several tick cell lines derived from both a natural vector (*Hyalomma anatolicum*) and other tick species not implicated in natural transmission of the virus (Bell-Sakyiet al., 2012).

The virus from an outbreak of “Crimean haemorrhagic fever” in the Crimean Peninsula in 1944 was not isolated or characterised until 1967. “Congo haemorrhagic fever” virus, isolated from a patient in the former Zaire (now Democratic Republic of the Congo) in 1956, was shown in 1969 to be the same virus. As a consequence, the names of both countries have been used in combination to describe the disease (Hoogstraal, 1979). Distribution of the virus reflects the broad distribution of *Hyalomma* ticks, the predominant vector of the virus (Avsic-Zupanc, 2007; Gard et al., 2011; Papa et al., 2011; Swanepoel & Paweska, 2011).

The natural cycle of CCHFV includes transovarial and transstadial transmission among ticks and a tick-vertebrate-tick cycle involving a variety of wild and domestic animals. Infection can also be transferred between infected and uninfected ticks during co-feeding on a host; so called ‘non-viraemic transmission’ phenomenon. *Hyalomma* ticks feed on a variety of domestic ruminants (sheep, goats, and cattle), and wild herbivores, hares, hedgehogs, and certain rodents. CCHFV infection in animals was reviewed by Nalca & Whitehouse (2007). Experimental infections of wild animals and livestock with CCHFV were reviewed by Spengler et al. (2016). Although animal infections are generally subclinical, the associated viraemia levels are sufficient to enable virus transmission to uninfected ticks (Swanepoel & Burt, 2004; Swanepoel & Paweska, 2011). Many birds are resistant to infection, but ostriches appear to be more susceptible than other bird species (Swanepoel et al., 1998). Although they do not appear to become viraemic, ground feeding birds may act as a vehicle for spread of CCHFV infected ticks. Results from serological surveys conducted in Africa and Eurasia indicate extensive circulation of the virus in livestock and wild vertebrates (Swanepoel & Burt, 2004).

Humans acquire infection from tick bites, or from contact with infected blood or tissues from livestock or human patients. After incubation humans can develop a severe disease with a prehaemorrhagic phase, a haemorrhagic phase, and a convalescence period (Hawman & Feldmann, 2023). Haemorrhagic manifestations can range from petechiae to large haematomas. Bleeding can be observed in the nose, gastrointestinal system, uterus and urinary tract, and the respiratory tract, with a case fatality rate ranging from 5% to 80% (Ergonul, 2006; Yen et al., 1985; Yilmaz et al., 2008). The severity of CCHF in humans highlights the impact of this zoonotic disease on public health. Although CCHFV has no economic impact on livestock animal production, the serological screening of animal serum samples for CCHFV-specific antibodies is very important. As seroprevalence in animals is a good indicator for local virus circulation, such investigations allow identification of high-risk areas for human infection (Mertens et al., 2013). Slaughterhouse workers, veterinarians, stockmen and others involved with the livestock industry should be made aware of the disease. They should take practical steps to limit or avoid exposure of naked skin to fresh blood and other animal tissues, and to avoid tick bites and handling ticks. Experiences from South Africa demonstrated that the use of repellents on animals before slaughter could reduce the numbers of infected slaughterhouse workers (Swanepoel et al., 1998). The treatment of livestock in general can reduce the tick density among these animals and thus reduce the risk of tick bite in animal handlers (Mertens et al., 2013). Such tick control by the use of acaricides is possible to some extent, but may be difficult to implement under extensive farming conditions. Inactivated mouse brain vaccine for the prevention of human infection has been used on a limited scale in Eastern Europe and the former USSR (Swanepoel & Paweska, 2011). Progress in CCHFV vaccine development is being made with several different approaches trialled to overcome current challenges (Dowall et al., 2017).

Infectivity of CCHFV is destroyed by boiling or autoclaving and low concentrations of formalin or beta-propiolactone. The virus is sensitive to lipid solvents. It is labile in infected tissues after death, presumably due to a fall in pH, but infectivity is retained for a few days at ambient temperature in serum, and for up to 3 weeks at 4°C. Infectivity is stable at temperatures below –60°C (Swanepoel & Paweska, 2011). CCHFV should be handled with appropriate biocontainment measures determined by risk analysis as described in Chapter 1.1.4 *Biosafety and biosecurity: Standard for managing biological risk in the veterinary laboratory and animal facilities* (Palmer, 2011; Whitehouse, 2004).

## B. DIAGNOSTIC TECHNIQUES

*Table 1. Diagnostic test formats for Crimean-Congo haemorrhagic fever virus infections in animals*

Method	Purpose					
	Population freedom from infection	Individual animal freedom from infection prior to movement	Contribute to eradication policies	Confirmation of clinical cases in animals	Prevalence of infection – surveillance	Immune status in individual animals or populations post-vaccination
Detection and identification of the agent <sup>(a)</sup>						
Real-time RT-PCR	–	++	–	+++ <sup>(b)</sup>	+ <sup>(c)</sup>	–
Virus isolation in cell culture	–	–	–	+ <sup>(b)</sup>	–	–
Detection of immune response						
IgG ELISA	+++	+	–	++ <sup>(d)</sup>	+++	–
Competitive ELISA	+++	+	–	++ <sup>(d)</sup>	+++	–
IgM ELISA	–	++	–	++ <sup>(e)</sup>	–	–

Key: +++ = recommended for this purpose; ++ recommended but has limitations; + = suitable in very limited circumstances; – = not appropriate for this purpose.

RT-PCR = reverse-transcription polymerase chain reaction; ELISA = enzyme-linked immunosorbent assay.

<sup>(a)</sup>A combination of agent identification methods applied on the same clinical sample is recommended.

<sup>(b)</sup>Molecular testing/isolation can be used to confirm acute infection in rare cases in animals showing clinical signs as viraemia tends to be transient.

<sup>(c)</sup>RT-PCR is used for the screening of tick populations in the context of surveillance studies.

<sup>(d)</sup>Serological evidence of active infection with CCHFV has been demonstrated by seroconversion based on a rise in total or IgG antibody titres on samples taken at 2–4 weeks apart.

<sup>(e)</sup>Serological evidence of active infection with CCHFV has been demonstrated by the detection of IgM antibodies specific to CCHFV using two different ELISAs based on two different antigens.

CCHFV infection causes only a mild fever in domestic and wild vertebrate animals with a detectable viraemia of up to 2 weeks (Gonzalez *et al.*, 1998; Gunes *et al.*, 2011). Similarly infected ostriches develop only low and short-lived viraemia and no clinical signs (Swanepoel & Burt, 2004). Therefore, recent infections in animals are rarely diagnosed and methods such as polymerase chain reaction (PCR), virus isolation in cell culture and IgM detection by enzyme-linked immunosorbent assay (ELISA) are mainly used in human CCHF diagnostics or in the special case that an animal has to be classified as CCHFV free. For prevalence analysis and for determination of whether CCHFV is circulating in a country, methods for the detection of IgG antibodies are preferred (Table 1). If there is any possibility or suspicion that diagnostic samples could be contaminated with CCHFV, they should be handled under an adequate biosafety level and all persons dealing with those samples should be aware of the possible risk and should use personal protective equipment to avoid human infections.

### 1. Detection and identification of the agent

For testing animals for viraemia, rapid diagnosis can be achieved by detection of viral nucleic acid in serum or plasma using conventional (Burt *et al.*, 1998) or real-time reverse transcription (RT-) PCR (Drosten *et al.*, 2002; Duh *et al.*, 2006; Koehler *et al.*, 2018; Negrodo *et al.*, 2017; Sas *et al.*, 2018; Wolfel *et al.*, 2007), or by demonstration of viral antigen (Shepherd *et al.*, 1988). Specimens to be submitted for laboratory confirmation of CCHF include blood and liver samples. Because of the risk of laboratory-acquired infections, work with CCHFV should be conducted in appropriate biosafety facilities.

The virus can be isolated from serum and organ suspensions in a wide variety of cell cultures, including Vero, LLC-MK2, SW-13, BSR-T7/5, CER and BHK21 cells, and identified by immunofluorescence using specific antibodies. Isolation and identification of virus can be achieved in 1–5 days, but cell cultures lack sensitivity and usually only detect high concentrations of virus present in the blood.

## 1.1. Virus isolation in cell culture

CCHFV can be isolated in mammalian cell cultures. Vero cells are commonly used, usually yielding an isolate between 1 and 5 days post-inoculation (p.i). CCHFV is poorly cytopathic and thus infectivity is titrated by demonstration of immunofluorescence in infected cells (Shepherd *et al.*, 1986). SW-13 cell line has also been used extensively for virus isolation, producing plaques within 4 days (p.i.). Identification of a CCHFV isolate has to be confirmed by immunofluorescence or molecular techniques (Burt *et al.*, 1998; Shepherd *et al.*, 1986).

### 1.1.1. Test procedure

- i) Susceptible cell lines include Vero-E6, BHK-21, LLC-MK2 and SW-13 cells. Inoculate 80% confluent monolayers of the preferred cell line with the specimen. The volume of specimen to be used depends on the size of the culture vessel (i.e. 25 cm<sup>2</sup> culture flask or 6- or 24-well tissue culture plate). The specimen volume should be sufficient to cover the cell monolayer. Samples of insufficient volumes can be diluted with tissue culture medium to prepare sufficient inoculation volume.
- ii) Adsorb the specimen for 1 hour at 37°C.
- iii) Remove inoculum. Add fresh tissue culture medium containing 2% fetal calf serum and other required additives, as per specific medium and cell line requirements.
- iv) Incubate at 37°C and 5% CO<sub>2</sub> for 4–7 days.
- v) Test supernatant for presence of CCHFV viral RNA using real-time RT-PCR as described below, or perform immunofluorescence assay on cell scrapings.
- vi) Isolates of CCHFV from clinical specimens cause no microscopically recognisable cytopathic effects (CPE) in most of these cell lines.

## 1.2. Nucleic acid detection

Molecular-based diagnostic assays, such as RT-PCR, serve as the front-line tool in the diagnosis of CCHF, as well as other viral haemorrhagic fevers (Drosten *et al.*, 2003). The benefit of molecular diagnostic assays is their rapidity compared to virus culture, often allowing a presumptive diagnosis to be reported within a few hours after receiving a specimen (Burt *et al.*, 1998). The RT-PCR is a sensitive method for diagnosis, but because of the genetic diversity of CCHFV, there might be some challenges with regard to design of primers or probes that allow detection of all circulating strains of the virus. Indeed, based on geographical origin and phylogenetic analyses of the S gene segment, CCHFV has previously been classified into nine geographical clades – four predominantly diffused in Africa, three in Europe, and two in Asia. Several real-time RT-PCR assays that detect strains from different geographical locations have been evaluated (Gruber *et al.*, 2019). While some assays have been shown to be highly sensitive, detecting as little as 10 viral RNA copies per ml of plasma, it is necessary to combine at least two molecular assays to ensure detection of the different CCHFV clades (Gruber *et al.*, 2019). The best assay combination(s) with the best detection efficacy for each CCHFV clade, based on all CCHFV sequences known at the time of the study, are shown in Table 2. In addition, a low-density microarray has been extensively validated in clinical specimens collected from confirmed cases of CCHF over 20 years by a WHO reference laboratory. It was shown to detect as few as 6.3 genome copies per reaction (Wolfel *et al.*, 2009).

**Table 2. Molecular assay combinations for the detection of CCHFV-specific nucleic acid**

Clade	Molecular assay combinations	Primer and probe names (5' → 3' sequence)
Africa 1	Real-time RT-PCR	Fwd CCRRealP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRRealP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)
Africa 2	Real-time RT-PCR	Fwd CCRRealP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRRealP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)
	Real-time RT-PCR	Fwd CCHF-SF2 (GGA-VTG-GTG-VAG-GGA-RTT-TG)

Clade	Molecular assay combinations	Primer and probe names (5' → 3' sequence)
		Rev CCHF-SR2 (CAD-GGT-GGR-TTG-AAR-GC) Fwd CCHF-N2 (CAA-RGG-CAA-RTA-CAT-MAT)
Africa 3	Nested RT-PCR	Fwd CCHF1 (CTG-CTC-TGG-TGG-AGG-CAA-CAA) Rev CCHF2_5 (TGG-GTT-GAA-GGC-CAT-GAT-GTA-T) Nested Fwd CCHFn15 (AGG-TTT-CCG-TGT-CAA-TGC-AAA) Nested Rev CCHFn25 (TTG-ACA-AAC-TCC-CTG-CAC-CAG-T)
	Nested RT-PCR	Fwd CrCon1+ (RWA-AYG-GRC-TTR-TGG-AYA-CYT-TCA-C) Rev CrCon1- (TRG-CAA-GRC-CKG-TWG-CRA-CWA-GWG-C) Nested Fwd CriCon2+ (ART-GGA-GRA-ARG-AYA-TWG-GYT-TYC-G) Nested Rev CriCon2- (CYT-TGA-YRA-AYT-CYC-TRC-ACC-ABT-C)
Africa 4	Real-time RT-PCR	Fwd CCRRealP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRRealP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)
	Real-time RT-PCR	Fwd CCHF-III (CAA-GAG-GTA-CCA-AGA-AAA-TGA-AGA-AGG-C) Rev CCHF-III-r (GCC-ACG-GGG-ATT-GTC-CCA-AAG-CAG-AC) Probe CCHFprobe-1 (ATC-TAC-ATG-CAC-CCT-GCY-GTG-YTG-ACA) Probe CCHFprobe-2 (TTC-TTC-CCC-CAC-TTC-ATT-GGR-GTG-CTC-A)
Asia 1	Nested PCR	Fwd CCF-115F (AAR-GGA-AAT-GGA-CTT-RTG-GA) Fwd CCF-131F (TGG-AYA-CYT-TCA-CAA-ACT-CC) Rev CCF-759R (GCA-AGG-CCT-GTW-GCR-ACA-AGT-GC)
	Real-time RT-PCR	Fwd CC1a_for (GTG-CCA-CTG-ATG-ATG-CAC-AAA-AGG-ATT-CCA-TCT) Rev CC1a_rev (GTG-CCA-CTG-ATG-ATG-CAC-AAA-AGG-ATT-CCA-TCT) Probe CCHF-01 (CAA-CAG-GCT-GCT-CTC-AAG-TGG-AG)
	Real-time RT-PCR	Fwd CCHF-SF2 (GGA-VTG-GTG-VAG-GGA-RTT-TG) Rev CCHF-SR2 (CAD-GGT-GGR-TTG-AAR-GC) Probe CCHF-N2 (CAA-RGG-CAA-RTA-CAT-MAT)
Asia 2	Nested PCR	Fwd CCF-115F (AAR-GGA-AAT-GGA-CTT-RTG-GA) Fwd CCF-131F (TGG-AYA-CYT-TCA-CAA-ACT-CC) Rev CCF-759R (GCA-AGG-CCT-GTW-GCR-ACA-AGT-GC)
	Real-time RT-PCR	Fwd (GAT-GAG-ATG-AAC-AAG-TGG-TTT-GAA-GA) Rev (GTA-GAT-GGA-ATC-CTT-TTG-TGC-ATC-AT)
	RT-PCR	Fwd CCS (ATG-CAG-GAA-CCA-TTA-ART-CTT-GGG-A) Rev 1 CCAS1 (CTA-ATC-ATA-TCT-GAC-AAC-ATT-TC) Rev 2 CCAS2 (CTA-ATC-ATG-TCT-GAC-AGC-ATC-TC)
Europe 1	Real-time RT-PCR	Fwd CCRRealP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRRealP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)
	Nested RT-PCR	Fwd CCF-115F (AAR-GGA-AAT-GGA-CTT-RTG-GA) Fwd CCF-131F (TGG-AYA-CYT-TCA-CAA-ACT-CC) Rev CCF-759R (GCA-AGG-CCT-GTW-GCR-ACA-AGT-GC)
Europe 2	Nested RT-PCR	Fwd CrCon1+ (RWA-AYG-GRC-TTR-TGG-AYA-CYT-TCA-C) Rev CrCon1- (TRG-CAA-GRC-CKG-TWG-CRA-CWA-GWG-C) Fwd CriCon2+ (ART-GGA-GRA-ARG-AYA-TWG-GYT-TYC-G) Rev CriCon2- (CYT-TGA-YRA-AYT-CYC-TRC-ACC-ABT-C)
Europe 3	Real-time RT-PCR	Fwd CCRRealP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRRealP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)

Clade	Molecular assay combinations	Primer and probe names (5' → 3' sequence)
	Real-time RT-PCR	Fwd CCRéalP1 (TCT-TYG-CHG-ATG-AYT-CHT-TYC) Rev CCRéalP2 (GGG-ATK-GTY-CCR-AAG-CA) Probe (ACA-SRA-TCT-AYA-TGC-AYC-CTG-C)
	Nested RT-PCR	Fwd CrCon1+ (RWA-AYG-GRC-TTR-TGG-AYA-CYT-TCA-C) Rev CrCon1- (TRG-CAA-GRC-CKG-TWG-CRA-CWA-GWG-C) Rev Fwd CriCon2+ (ART-GGA-GRA-ARG-AYA-TWG-GYT-TYC-G) Nested Rev CriCon2- (CYT-TGA-YRA-AYT-CYC-TRC-ACC-ABT-C)
All	Real-time RT-PCR	Fwd CCHF-SF2 (GGA-VTG-GTG-VAG-GGA-RTT-TG) Rev CCHF-SR2 (CAD-GGT-GGR-TTG-AAR-GC) Probe CCHF-N2 (CAA-RGG-CAA-RTA-CAT-MAT)
	RT-PCR	Fwd CCS (ATG-CAG-GAA-CCA-TTA-ART-CTT-GGG-A) Rev 1 CCAS1 (CTA-ATC-ATA-TCT-GAC-AAC-ATT-TC) Rev 2 CCAS2 (CTA-ATC-ATG-TCT-GAC-AGC-ATC-TC)
	Real-time RT-PCR	Fwd CC1a_for (GTG-CCA-CTG-ATG-ATG-CAC-AAA-AGG-ATT-CCA-TCT) Rev CC1a_rev (GTG-CCA-CTG-ATG-ATG-CAC-AAA-AGG-ATT-CCA-TCT) Probe CCHF-01 (CAA-CAG-GCT-GCT-CTC-AAG-TGG-AG)

(Data and table modified from Gruber *et al.* 2019)

## 2. Serological tests

Virus neutralisation assays, generally considered to be highly specific, are rarely used for CCHFV diagnosis. Members of the *Orthonairovirus* genus generally induce a weaker neutralising antibody response than members of other genera in the family *Nairoviridae*. Another drawback is the necessity to perform this assay in high biosafety containment because it uses live virus (Burt *et al.*, 1994; Rodriguez *et al.*, 1997).

Currently, there are only a few CCHFV commercial kits for IgM or IgG by ELISA or immunofluorescence (IFA). These are all designed for the human diagnostic market. However, it is possible to adapt these commercial ELISAs and IFAs for serological testing in animals. In addition, some in-house ELISAs have been published for the detection of CCHFV-specific antibodies in animals.

Diagnostic performance for humans have been compared between the methods using sensitivity, specificity, concordance and degree of agreement with particular focus on the phase of the infection (Emmerich *et al.*, 2021). Available serological test systems detect anti-CCHFV IgM and IgG antibodies accurately, but their diagnostic performance varies with respect to the phase of the infection. In the early and convalescent phases of infection, the sensitivity for detecting specific IgG antibodies differed for the ELISA. Both test systems based on immunofluorescence showed an identical sensitivity for detection of anti-CCHFV IgM antibodies in acute and convalescent phases of infection.

IgM antibodies in livestock (sheep, goat and cattle) can be detected by using an IgM-capture ELISA. IgG antibodies can be detected by an IgG-sandwich or indirect ELISA, and total antibodies can be detected by competition ELISA. The benefit of competitive ELISA is the capacity to investigate different animal species, because they are host species independent. Commercial kits for the detection of CCHFV-specific antibodies or the detection of viral antigen are available. The limiting factor for the replication of these protocols in other laboratories is the availability of antigens and (where relevant) specified monoclonal antibodies. Most of the tests described for livestock and wild animals have not undergone a formal validation process (Mertens *et al.*, 2013). One of the biggest challenges for such validation studies is the availability of an adequate number of positive well characterised control samples.

For information on the availability of reference reagents for use in veterinary diagnostic laboratories, contact the WOA Collaborating Centres for Zoonoses in Europe and in Asia-Pacific.

## C. REQUIREMENTS FOR VACCINES

There is no vaccine available for animals.

## REFERENCES

- AVSIC-ZUPANC T. (2007). Epidemiology of Crimean–Congo hemorrhagic fever in the Balkans. *In: Crimean–Congo Hemorrhagic Fever, a Global Perspective*, Ergonul O. & Whitehouse C.A., eds. Springer: Dordrecht, Netherlands, 75–88.
- BELL-SAKYI L., KOHL D., BENTE D.A. & FAZAKERLEY J.F. (2012). Tick cell lines for study of Crimean–Congo hemorrhagic fever virus and other arboviruses. *Vector Borne Zoonotic Dis.*, **12**, 769–781.
- BURT F.J., LEMAN P.A., ABBOTT J.C. & SWANEPOEL R. (1994). Serodiagnosis of Crimean–Congo haemorrhagic fever. *Epidemiol. Infect.*, **113**, 551–562. Doi: 10.1017/s0950268800068576
- BURT F.J., LEMAN P.A., SMITH J.F. & SWANEPOEL R. (1998). The use of a reverse transcription-polymerase chain reaction for the detection of viral nucleic acid in the diagnosis of Crimean–Congo haemorrhagic fever. *J. Virol. Methods*, **70**, 129–37.
- DROSTEN C., GOTTING S., SCHILLING S., ASPER M., PANNING M., SCMITZ H. & GUNTER S. (2002). Rapid detection and quantification of RNA of Ebola and Marburg viruses, Lassa virus, Crimean–Congo hemorrhagic fever virus, Rift Valley fever virus, dengue virus, and yellow fever virus by real-time transcription-PCR. *J. Clin. Microbiol.*, **40**, 2323–2340.
- DROSTEN C., KUMMERER B.M., SCMITZ H. & GUNTER S. (2003). Molecular diagnosis of viral hemorrhagic fevers. *Antiviral Res.*, **57**, 61–87.
- DUH D., SAKSIDA A., PETROVEC M., DEDUSHAJ I. & AVSIC-ZUPANC T. (2006). Novel one-step real-time RT-PCR assay for rapid and specific diagnosis of Crimean–Congo hemorrhagic fever encountered in the Balkans. *J. Virol. Methods*, **133**, 175–179.
- EMMERICH P., MIKA A., VON POSSEL R., RACKOW A., LIU Y., SCHMITZ H., GÜNTHER S., SHERIFI K., HALILI B., JAKUPI X., BERISHA L., AHMETI S. & DESCHERMEIER C. (2018). Sensitive and specific detection of Crimean–Congo Hemorrhagic Fever Virus (CCHFV)-Specific IgM and IgG antibodies in human sera using recombinant CCHFV nucleoprotein as antigen in  $\mu$ -capture and IgG immune complex (IC) ELISA tests. *PLoS Negl. Trop. Dis.*, **12**(3):e0006366. doi: 10.1371/journal.pntd.0006366.
- EMMERICH P., VON POSSEL R., DESCHERMEIER C., AHMETI S., BERISHA L., HALILI B., JAKUPI X., SHERIFI K., MESSING C. & BORCHARDT-LOHÖLTER V. (2021). Comparison of diagnostic performances of ten different immunoassays detecting anti-CCHFV IgM and IgG antibodies from acute to subsided phases of Crimean–Congo hemorrhagic fever. *PLoS Negl. Trop. Dis.*, **15** (3):e0009280.
- ERGONUL O. (2006). Crimean–Congo haemorrhagic fever. *Lancet Infect. Dis.*, **6**, 203–214.
- GONZALEZ J.-P., CAMICAS J.-L., COMET J.-P. & WILSON M.L. (1998). Biological and clinical responses of West African sheep to Crimean–Congo haemorrhagic fever virus experimental infection. *Res. Virol.*, **149**, 445–455.
- GUNES T., POYRAZ O., VATANSEVER Z. (2011). Crimean–Congo hemorrhagic fever virus in ticks collected from humans, livestock, and picnic sites in the hyperendemic region of Turkey. *Vector Borne Zoonotic Dis.*, **11**, 1411–1416.
- GRARD G., DREXLER J.F., FAIR J., MUYEMBE J.-J., WOLFE N.D., DROSTEN C. & LEROY E.M. (2011). Re-emergence of Crimean–Congo hemorrhagic fever virus in Central Africa. *PLoS Negl. Trop. Dis.*, **5**(10): e1350. doi:10.1371/journal.pntd.0001350.
- GRUBER C., BARTOLINI B., CASTILLETI C., MIRAZIMI A., HEWSON R., CHRISTOVA I., AVŠIČ T., GRUNOW R., PAPA A., SÁNCHEZ-SECO M. P., KOPMANS M., IPPOLITO G., CAPOBIANCHI M. R., REUSKEN C. & DI CARO A. (2019). Geographical Variability Affects CCHFV Detection by RT-PCR: A Tool for *In-Silico* Evaluation of Molecular Assays. *Viruses*, **11**, 953.
- GULCE-İZ S., ELALDI N., CAN H Şahar E.A., Karakavuk M., Gül A., Kumoğlu G.Ö., Döşkaya A.D., Gürüz A.Y., Özdarendeli A., Felgner P.L., Davies H. & Döşkaya M. (2021). Development of a novel recombinant ELISA for the detection of Crimean–Congo hemorrhagic fever virus IgG antibodies. *Sci. Rep.*, **11**, 5936. doi:10.1038/s41598-021-85323-1.

- HAWMAN D.W. & FELDMANN H. (2023). Crimean–Congo haemorrhagic fever virus. *Nat. Rev. Microbiol.*, **21**, 463–477 <https://doi.org/10.1038/s41579-023-00871-9>
- HOOGSTRAAL H. (1979). The epidemiology of tick-borne Crimean–Congo haemorrhagic fever in Asia, Europe and Africa. *J. Med. Entomol.*, **15**, 307–417.
- KOEHLER J.W., DELP K.L., HALL A.T., OLSCHNER S.P., KEARNEY B.J., GARRISON A.R., ALTAMURA L.A., ROSSI C.A. & MINOGUE T.D. (2018). Sequence Optimized Real-Time Reverse Transcription Polymerase Chain Reaction Assay for Detection of Crimean–Congo Hemorrhagic Fever Virus. *Am. J. Trop. Med. Hyg.*, **98**, 211–215.
- MERTENS M., SCHMIDT K., OZKUL A. & GROSCHUP M.H. (2013). The impact of Crimean–Congo hemorrhagic fever virus on public health. *Antiviral Res.*, **98**, 248–260.
- NALCA A. & WHITEHOUSE C.A. (2007). Crimean–Congo hemorrhagic fever virus infection among animals. In: Crimean–Congo Hemorrhagic Fever: A Global Perspective, Ergonul O. & Whitehouse C.A., eds. Springer: Dordrecht, Netherlands, 155–165.
- NEGREDO A., DE LA CALLE-PRieto F., PALENCIA-HERREJÓN E., MORA-RILLO M., ASTRAY-MOCHALES J., SÁNCHEZ-SECO M.P., BERMEJO LOPEZ E., MENÁRGUEZ J., FERNÁNDEZ-CRUZ A., SÁNCHEZ-ARTOLA B., KEOUGH-DELGADO E., RAMÍREZ DE ARELLANO E., LASALA F., MILLA J., FRAILE J.L., ORDOBÁS GAVÍN M., MARTINEZ DE LA GÁNDARA A., LÓPEZ PEREZ L., DIAZ-DIAZ D., LÓPEZ-GARCÍA M.A., DELGADO-JIMENEZ P., MARTÍN-QUIRÓS A., TRIGO E., FIGUEIRA J.C., MANZANARES J., RODRIGUEZ-BAENA E., GARCIA-COMAS L., RODRÍGUEZ-FRAGA O., GARCÍA-ARENZANA N., FERNÁNDEZ-DÍAZ M.V., CORNEJO V.M., EMMERICH P., SCHMIDT-CHANASIT J., ARRIBAS J.R., CRIMEAN CONGO HEMORRHAGIC FEVER@MADRID WORKING GROUP (2017). Autochthonous Crimean–Congo Hemorrhagic Fever in Spain. *N. Engl. J. Med.*, **377**, 154–161. 10.1056/NEJMoa1615162.
- PALMER S. (2011). Deliberate release of zoonotic agents. In: Oxford Textbook of Zoonosis: Biology, Clinical Practise and Public Health Control, Second Edition, Palmer S.R., Soulsby L., Torgerson P.R. & Brown D.W.G., eds. Oxford University Press, UK, p. 1214.
- PAPA A., TZALA E. & MALTEZOU H. (2011). Crimean–Congo hemorrhagic fever virus, Northeastern Greece. *Emerg. Infect. Dis.*, **17**, 141–143.
- RODRIGUEZ L.L., MAUPIN G.O., KSIAZEK T.G., ROLLIN P.E., KHAN A.S., SCHWARZ T.F., LOFTS R.S., SMITH J.F., NOOR A.M., PETERS C.J. & NICHOL S.T. (1997). Molecular investigation of a multisource outbreak of Crimean–Congo hemorrhagic fever in the United Arab Emirates. *Am. J. Trop. Med. Hyg.*, **57**, 512–518.
- SAS M.A., VINA-RODRIGUEZ A., MERTENS M., EIDEN M., EMMERICH P., CHAINTOUTIS S.C., MIRAZIMI A. & GROSCHUP M.H. (2018). A one-step multiplex real-time RT-PCR for the universal detection of all currently known CCHFV genotypes. *J. Virol. Methods*, **255**, 38–43.
- SHEPHERD A.J., SWANEPOEL R. & GILL D.E. (1988). Evaluation of enzyme-linked immunosorbent assay and reversed passive hemagglutination for detection of Crimean–Congo hemorrhagic fever virus antigen. *J. Clin. Microbiol.*, **26**, 347–353.
- SHEPHERD A.J., SWANEPOEL R., LEMAN P.A. & SHEPHERD S.P. (1986). Comparison of methods for isolation and titration of Crimean-Congo hemorrhagic fever virus. *J. Clin. Microbiol.*, **24**, 654–656.
- SPENGLER J.R., ESTRADA-PEÑA A., GARRISON A.R., SCHMALJOHN C., SPIROPOULOU C.F., BERGERON É. & BENTE D.A. (2016). A chronological review of experimental infection studies of the role of wild animals and livestock in the maintenance and transmission of Crimean-Congo hemorrhagic fever virus. *Antiviral Res.*, **135**, 31–47. doi: 10.1016/j.antiviral.2016.09.013.
- SWANEPOEL R & BURT F.J. (2004). Crimean–Congo haemorrhagic fever. Second Edition. In: Infectious diseases of livestock with special reference to South Africa, Coetzer J.A.W, Tustin R.C., eds. Cape Town: Oxford University Press Southern Africa, pp. 1077–1085.
- SWANEPOEL R., LEMAN P.A., BURT, F.J., JARDINE J., VERWOERD D.J., CAPUA I., BRUCKNER G.K. & BURGER W.P. (1998). Experimental infection of ostriches with Crimean-Congo haemorrhagic fever virus. *Epidemiol. Infect.*, **121**, 427–432.



SWANEPOEL R. & PAWESKA J.T. (2011). Crimean-Congo hemorrhagic fever. *In: Oxford Textbook of Zoonosis: Biology, Clinical Practise and Public Health Control*, Second Edition. Palmer S.R., Soulsby L., Torgerson P.R. & Brown D.W.G.,eds. Oxford University Press, UK, pp. 287–293.

WHITEHOUSE C.A. (2004). Crimean–Congo hemorrhagic fever. *Antivir. Res.*, **64**, 145–160.

WOLFEL R., PAWESKA J.T., PETERSEN N., GROBBELAAR A.G., LEMAN P.A., HEWSON R., GEORGES-COURBOT, M., PAPA, A., GÜNTER S. & DROSTEN C. (2007). Virus detection and monitoring of viral load in Crimean-Congo hemorrhagic fever virus patients. *Emerg. Infect. Dis.*, **13**, 1097–1100.

WOLFEL R., PAWESKA J.T., PETERSEN N., GROBBELAAR A.G., LEMAN P.A., HEWSON R., GEORGES-COURBOT M., PAPA, A., HEISER V., PANNING M., GUNTER S. & DROSTEN C. (2009). Low-density microarray for rapid detection and identification of Crimean–Congo hemorrhagic fever virus. *J. Clin. Microbiol.*, **47**, 1025–1030.

YEN Y.C., KONG L.X., LEE L., ZHANG Y.Q., LI F., CAI B.J. & GAO S.Y. (1985). Characteristics of Crimean–Congo hemorrhagic fever virus (Xinjiang strain) in China. *Am. J. Trop. Med. Hyg.*, **34**, 1179–1182.

YILMAZ G.R., BUZGAN T., TORUNOGLU M.A., SAFRAN A., IRMAK H., COM S., UYAR Y., CARHAN A., OZKAYA E. & ERTEK M. (2008). A preliminary report on Crimean–Congo haemorrhagic fever in Turkey, March–June 2008. *Euro Surveill.*, **13**.

\*  
\* \*

**NB:** At the time of publication (2024) there was no WOA Reference Laboratory for Crimean–Congo haemorrhagic fever (please consult the WOA Web site: <https://www.woah.org/en/what-we-offer/expertise-network/reference-laboratories/#ui-id-3>).

**NB:** FIRST ADOPTED IN 2014. MOST RECENT UPDATES ADOPTED 2024.